

# Single eta production in heavy quarkonia: breakdown of multipole expansion

Yu.A.Simonov, A.I.Veselov

State Research Center

Institute of Theoretical and Experimental Physics,  
Moscow, 117218 Russia

December 26, 2008

## Abstract

The  $\eta$  production in the  $(n, n')$  bottomonium transitions  $\Upsilon(n) \rightarrow \Upsilon(n')\eta$ , is studied in the method used before for dipion heavy quarkonia transitions. The widths  $\Gamma_\eta(n, n')$  are calculated without fitting parameters for  $n = 2, 3, 4, 5, n' = 1$ . Resulting  $\Gamma_\eta(4, 1)$  is found to be large in agreement with recent data. Multipole expansion method is shown to be inadequate for large size systems considered.

## 1 Introduction

The  $\eta$  and  $\pi^0$  production in heavy quarkonia transitions is attracting attention of experimentalists for a long time [1]. The first result refers to the  $\psi(2S) \rightarrow J/\psi(1S)\eta$  process (to be denoted as  $\psi(2, 1)\eta$  in what follows, similarly for  $\Upsilon$ ) with  $\frac{\Gamma_\eta}{\Gamma_{tot}} = (3.09 \pm 0.08)\%$  [1],  $\Gamma_{tot} = 337 \pm 13$  keV.

For the  $\Upsilon(2, 1)\eta$  and  $\Upsilon(3, 1)\eta$  transitions only upper limits  $B < 2 \cdot 10^{-3}$  and  $B < 2.2 \cdot 10^{-3}$  were obtained in [2] and [3] correspondingly and preliminary results appeared recently in [4],  $B(\Upsilon(2, 1)\eta) = (2.5 \pm 0.7 \pm 0.5)10^{-4}$ ,  $B(\Upsilon(2, 1)\pi^0) < 2.1 \cdot 10^{-4}(90\% \text{ c.l.})$  and  $B(\Upsilon(3, 1)\eta) < 2.9 \cdot 10^{-4}$  in [5].

On theoretical side the dominant approach for both dipion and single  $\eta$  and  $\pi$  production is the Multipole Expansion Method (MEM) (see [6, 7] and refs. therein), where it is assumed that heavy quarks emit two gluons and the

latter are converted into meson(s) by a not clarified mechanism. An essential requirement for this mechanism in QED is the smallness of the source size  $r_0$  as compared to the wavelength, so that  $r_0 k \ll 1$ .

In reality both for charmonium and bottomonium transitions  $r_0 k \gtrsim 1$ , but it is not this parameter which invalidates MEM for heavy quarkonia. It appears, that in QCD there is another important length parameter, the QCD vacuum correlation length  $\lambda$ , which makes it impossible to emit freely gluons at points separated by distance  $r, r > \lambda$ .

The value of  $\lambda$  was found on the lattice and analytically,  $\lambda \lesssim 0.2$  fm [8]. Since r.m.s. radii of all excited  $c\bar{c}, b\bar{b}$  states are larger than  $0.5$  fm<sup>1</sup> all vacuum gluons there are correlated forming the QCD string and emission of additional gluons (if any) implies formation of heavy hybrids. All this is considered in detail in Field Correlator Method (FCM) [9].

One can make an independent check of MEM in application to the bottomonium level calculation. Here MEM yields nonperturbative correction to the levels expressed via gluonic condensate [10]. Comparison to the experimental data shows (see [11] and Table 1 below) that for all level splittings except  $(2S - 1S)$  in bottomonium, MEM prediction is more than 50% off, while for charmonium MEM does not work at all. Thus one concludes that only at distances below or equal  $0.2$  fm, MEM can give reasonable results, while for all states of charmonium and all excited states of bottomonium (where sizes are much larger than vacuum correlation length  $\lambda$ ) the application of MEM is unjustifiable.

**Table 1.** Predicted splittings, between spin-averaged levels, and experiment.

| <i>splitting</i>     | <i>MEM</i> [11] | <i>FCM</i> [12] | <i>exp.</i> [1] |
|----------------------|-----------------|-----------------|-----------------|
| $2S - 1S (b\bar{b})$ | 479             | 557             | 558 MeV         |
| $2S - 1P (b\bar{b})$ | 181             | 122             | 123 MeV         |
| $3S - 2S (b\bar{b})$ | 4 570           | 348             | 332 MeV         |
| $2S - 1S (c\bar{c})$ | 9 733           | 610             | 606 MeV         |

---

<sup>1</sup>In reality the r.m.s. radius  $r_0$  of heavy quarkonia is not small, e.g. for  $\Upsilon(nS)$  it is equal to  $0.2$  fm,  $0.5$  fm,  $0.7$  fm,  $0.9$  fm, for  $n = 1, 2, 3, 4, 5$ , and for charmonium this radius is even larger:  $0.4$  fm and  $0.8$  fm for  $n = 1$  and  $2$  respectively.

A similar failure of MEM is found in applications to dipion bottomonium transitions, where using MEM one can fit dipion spectra in  $\Upsilon(2,1)\pi\pi$ , but not in  $\Upsilon(3,1)\pi\pi$  and  $\Upsilon(4,2)\pi\pi$  [6, 7]. In contrast to that, FCM as will be discussed below explains both spectra and  $\cos\theta$  dependence for all dipion transitions in universal approach with two fixed parameters.

In FCM large distances are under control and not single gluons but combined effect of all gluons in the string defines the dynamics.

In particular, single eta emission in heavy quarkonia proceeds via string breaking due to  $q\bar{q}$  pair creation with simultaneous emission of  $\pi$  or  $\eta$ . The flavor SU(3) violation in  $\eta$  production then resides in difference of threshold positions and wave functions for  $B\bar{B}$  and  $B_s\bar{B}_s$  ( $D\bar{D}$  and  $D_s\bar{D}_s$ ) intermediate states.

As it is clear, the dynamics of FCM for  $\eta$  emission does not depend strongly on heavy quark mass, and only sizes of initial and final heavy quarkonia states and intermediate heavy-light mesons enter in the form of overlap matrix elements.

In contrast to that, MEM predicts a strong dependence on the heavy quark mass,  $\Gamma_\eta(n, n') = O(\frac{1}{m_Q^2})$ . In addition in [7] a strong suppression of the ratio  $\Gamma_\eta/\Gamma_{\pi\pi}$  with the growth of the energy release  $\Delta M = M(n) - M(n')$ ,  $\Gamma_\eta/\Gamma_{\pi\pi} \sim \frac{p_\eta^3}{(\Delta M)^7}$  is predicted for higher excited states of quarkonia and bottomonia, which does not agree with experiment (see below).

Using models based on MEM in [6] small ratios of widths

$$\frac{\Gamma(\Upsilon(2,1)\eta)}{\Gamma(\psi(2,1)\eta)} \cong 2.5 \cdot 10^{-3} \quad \text{and} \quad \frac{\Gamma(\Upsilon(3,1)\eta)}{\Gamma(\psi(2,1)\eta)} = 1.3 \cdot 10^{-3} \quad (1)$$

have been predicted, with the model property that the bottomonium yields of  $\eta$  would be smaller than those of charmonium; specifically in the method of [6], the width is proportional to  $p^3/m_Q^2$ , so that for  $\Upsilon(4,1)\eta$  the ratio  $\frac{\Gamma(\Upsilon(4,1)\eta)}{\Gamma(\psi(2,1)\eta)} \approx 3.3 \cdot 10^{-3}$ .

However recently [13] new BaBar data have been published on  $\Upsilon(4,1)\eta$  with the branching ratio

$$B(\Upsilon(4,1)\eta) = (1.96 \pm 0.06 \pm 0.09)10^{-4} \quad (2)$$

and [13]

$$\frac{\Gamma(\Upsilon(4,1)\eta)}{\Gamma(\Upsilon(4,1)\pi^+\pi^-)} = 2.41 \pm 0.40 \pm 0.12. \quad (3)$$

This latter result is very large, indeed the corresponding ratio  $\frac{\Gamma(\Upsilon(4,1)\eta)}{\Gamma(\psi(2,1)\eta)}$  is  $\approx 0.4$  and theoretical estimates (1) from [6] for a similar ratio yields  $3.3 \cdot 10^{-3}$ . Thus, the experimental ratio is very large as compared to MEM predictions [6, 7]. All this suggests that another mechanism can be at work in single  $\eta$  production and below we exploit the approach based on the Field Correlator Method (FCM) recently applied to  $\Upsilon(n, n')\pi\pi$  transitions with  $n \leq 3$  in [14, 15],  $n \leq 4$  in [16] and  $n = 5$  in [17, 18].

In this paper we confront MEM and FCM and show that recent experimental data on single  $\eta$  production in  $\Upsilon(4S) - \Upsilon(1S)$  transition give a strong support to the FCM result and cannot be explained in the framework of MEM.

The method essentially exploits the mechanism of Internal Loop Radiation (ILR) with light quark loop inside heavy quarkonium and has two fundamental parameters – mass vertices in chiral light quark pair  $q\bar{q}$  creation  $M_{br} \approx f_\pi$  and pair creation vertex without pseudoscalars,  $M_\omega \approx 2\omega$ , where  $\omega(\omega_s)$  is the average energy of the light (strange) quark in the  $B(B_s)$  meson. Those are calculated with relativistic Hamiltonian [12] and considered as fixed for all types of transitions  $\omega = 0.587$  GeV,  $\omega_s = 0.639$  GeV (see Appendix 1 of [14] for details).

Any process of heavy quarkonium transition with emission of any number of Nambu-Goldstone (NG) mesons is considered in ILR as proceeding via intermediate states of  $B\bar{B}, B\bar{B}^* + c.c., B_s\bar{B}_s$  etc. (or equivalently  $D\bar{D}$  etc.) with NG mesons emitted at vertices.

For one  $\eta$  or  $\pi^0$  emission one has diagrams shown in Fig.1, where dashed line is for the NG meson. As shown in [14, 15, 16], based on the chiral Lagrangian derived in [19], the meson emission vertex has the structure

$$\mathcal{L}_{CDL} = -i \int d^4x \bar{\psi}(x) M_{br} \hat{U}(x) \psi(x) \quad (4)$$

$$\hat{U} = \exp \left( i\gamma_5 \frac{\varphi_a \lambda_a}{f_\pi} \right), \varphi_a \lambda_a = \sqrt{2} \begin{pmatrix} \frac{\eta}{\sqrt{6}} + \frac{\pi^0}{\sqrt{2}}, & \pi^+, & K^+ \\ \pi^-, & \frac{\eta}{\sqrt{6}} - \frac{\pi^0}{\sqrt{2}}, & K^0 \\ K^-, & \bar{K}^0, & -\frac{2\eta}{\sqrt{6}} \end{pmatrix}, \quad (5)$$




Fig.1 Single eta production (dashed line) from  $\Upsilon(n)BB^*$  vertex (a), and  $BB^*\Upsilon(n')$  vertex (b).

The lines (1,2,3) in the  $\hat{U}$  matrix (2) refer to  $u, d, s$  quarks and hence to the channels  $B^+B^-$ ,  $B^0\bar{B}^0$ ,  $B_s^0\bar{B}_s^0$  (and to the corresponding channels with  $B^*$  instead of  $B$ ). Therefore the emission of a single  $\eta$  in heavy quarkonia transitions requires the flavour  $SU(3)$  violation and resides in our approach in the difference of channel contribution  $B\bar{B}^*$  and  $B_s\bar{B}_s^*$ , while the  $\pi^0$  emission is due the difference of  $B^0\bar{B}^{0*}$  and  $B^+B^{-*}$  channels (with  $B \rightarrow D$  for charmonia).

The paper is devoted to the explicit calculation of single  $\eta$  emission widths in bottomonium  $\Upsilon(n,1)\eta$  transitions with  $n = 2, 3, 4, 5$ . Since theory has no fitting parameters (the only ones,  $M_\omega$  and  $M_{br}$  are fixed by dipion transitions) our predictions depend only on the overlap matrix elements, containing wave functions of  $\Upsilon(nS)$ ,  $B, B_s, B^*, B_s^*$ . The latter have been computed previously in relativistic Hamiltonian technic in [12] and used extensively in dipion transitions in [16, 17, 18].

The paper is organized as follows. In section 2 general expressions for process amplitudes are given; in section 3 results of calculations are presented and discussed and a short summary and perspectives are given.

## 2 General formalism

The process of single NG boson emission in bottomonium transition is described by two diagraman depicted in Fig.1, (a) and (b) which can be written according to the general formalism of FCM [14, 16, 17] as (we consider  $\eta$  emission), see Appendix for more detail,

$$\mathcal{M} = \mathcal{M}_\eta^{(1)} + \mathcal{M}_\eta^{(2)}; \mathcal{M}_\eta^{(i)} = \mathcal{M}_{B_s B_s^*}^{(i)} - \mathcal{M}_{B B^*}^{(i)}, i = 1, 2 \quad (6)$$

For the diagram of Fig. 1(a) the amplitude for intermediate  $BB^*$  or  $B_s B_s^*$  state can be written as

$$\mathcal{M}_\eta^{(1)} = \int \frac{J_n^{(1)}(\mathbf{p}, \mathbf{k}) J_{n'}(\mathbf{p})}{E - E(\mathbf{p})} \frac{d^3 \mathbf{p}}{(2\pi)^3}, \quad (7)$$

while  $\mathcal{M}_\eta^{(2)}$ , corresponding to the diagram of Fig.1(b), has the same form, but without NG boson energy in the denominator of (7). The overlap integrals of  $\Upsilon(nS)$  and  $BB^*$  wave functions with emission of  $\eta$  with momentum  $\mathbf{k}$  are denoted by  $J_n^{(i)}(\mathbf{p}, \mathbf{k})$ , the corresponding integrals without  $\eta$  emission are given by  $J_{n'}(\mathbf{p})$ .

Finally we define all quantities in the denominator of (7); in  $\mathcal{M}_{BB^*}^{(1)}$  the denominator is

$$E - E(\mathbf{p}) = M(\Upsilon(nS)) - (\omega_\eta + M_B + M_B^* + \frac{\mathbf{p}^2}{2M_B} + \frac{(\mathbf{p} - \mathbf{k})^2}{2M_B^*}) \equiv -\Delta M^* - \omega_\eta - E(\mathbf{p}, \mathbf{k}). \quad (8)$$

For  $\mathcal{M}_\eta^{(2)}$  one omits  $\omega_\eta$  and  $\mathbf{k}$  in (8). Finally after taking Dirac traces in amplitudes and accounting for the  $p$ -wave of emitted  $\eta$  one can represent the matrix element  $\mathcal{M}_\eta^{(i)}$  as follows:

$$\mathcal{M}_\eta^{(1)} = e_{i'i} k_l \left( \frac{1}{\omega_s^3} \mathcal{L}_s^{(1)} - \frac{1}{\omega^3} \mathcal{L}^{(1)} \right) \quad (9)$$

Indices  $i'i$  in  $e_{i'i}$  in (9) refer to the  $\Upsilon(n'S)$  and  $\Upsilon(nS)$  polarizations respectively.

$$\mathcal{M}_\eta^{(2)} = e_{i'i} k_l \left( \frac{1}{\omega_s^3} \mathcal{L}_s^{(2)} - \frac{1}{\omega^3} \mathcal{L}^{(2)} \right) \quad (10)$$

with  $\mathcal{L}^{(i)}, \mathcal{L}_s^{(i)}$  being integrals of overlap factors  $J_n(\mathbf{p}, \mathbf{k}) J_{n'}(\mathbf{p})$ .

The width of the  $\Upsilon(n, n')\eta$  decay is obtained from  $|\mathcal{M}|^2$  averaging over vector polarizations as

$$\Gamma_\eta = \frac{1}{3} \sum_{i,i'} |\mathcal{M}|^2 d\Phi = \frac{2k^2}{3} d\Phi \left| \left( \frac{\mathcal{L}_s^{(1)}}{\omega_s^3} - \frac{\mathcal{L}^{(1)}}{\omega^3} \right) + \left( \frac{\mathcal{L}_s^{(2)}}{\omega_s^3} - \frac{\mathcal{L}^{(2)}}{\omega^3} \right) \right|^2 \quad (11)$$

where the phase space factor  $d\Phi = \frac{d^3 k}{(2\pi)^3} 2\pi \delta(M(\Upsilon(n)) - M(\Upsilon(n')) - \omega_\eta - \frac{k^2}{2M(\Upsilon(n'))})$ .

Introducing the average  $\bar{\omega} = \frac{1}{2}(\omega_s + \omega)$ , one can rewrite (11) as

$$\Gamma_\eta = \left( \frac{M_{br}}{f_\eta} \right)^2 \left( \frac{M_\omega}{2\bar{\omega}} \right)^2 \zeta_\eta \frac{k^3}{\bar{\omega}^4} e^{-\frac{k^2}{2\beta_2^2}} \left| \left( \frac{\bar{\omega}}{\omega_s} \right)^3 \mathcal{L}_s^{(1)} - \left( \frac{\bar{\omega}}{\omega} \right)^3 \mathcal{L}^{(1)} + \left( \frac{\bar{\omega}}{\omega_s} \right)^3 \mathcal{L}_s^{(2)} - \left( \frac{\bar{\omega}}{\omega} \right)^3 \mathcal{L}^{(2)} \right|^2 \quad (12)$$

with  $\zeta_\eta = \frac{16}{9\pi N_c^2} \cong 0.063$ .

One can see from the general structure of  $\Gamma_\eta$ , that the main effect comes from the difference  $\left| \left( \frac{\bar{\omega}}{\omega_s} \right)^3 - \left( \frac{\bar{\omega}}{\omega} \right)^3 \right| \approx |0.882 - 1.139| \approx 0.257$ , and from the difference of  $|\mathcal{L}_s^{(i)} - \mathcal{L}^{(i)}| \leq 0.05$ .

### 3 Results and discussion

We consider here the single  $\eta$  emission in bottomonium transitions  $\Upsilon(n, 1)\eta$  with  $n = 2, 3, 4, 5$ . The corresponding values of  $\Delta M^*$ ,  $\Delta M_s^*$ ,  $\omega_\eta$ ,  $k$  are given in the Table 2.

**Table 2.** Mass parameters of  $\Upsilon(n, n')\eta$  transitions (all in GeV,  $k$  in GeV/c).

| $(n, n')$      | 2,1   | 3,1   | 4,1   | 5,1   |
|----------------|-------|-------|-------|-------|
| $\Delta M^*$   | 0.582 | 0.25  | 0.026 | -0.26 |
| $\Delta M_s^*$ | 0.757 | 0.425 | 0.20  | -0.08 |
| $\omega_\eta$  | 0.562 | 0.87  | 1.075 | 1.325 |
| $k$            | 0.115 | 0.674 | 0.923 | 1.20  |

The resulting values of  $\Gamma_\eta(n, n')$  have been computed as in (12) with  $\omega = 0.587$  GeV and  $\omega_s = 0.639$  GeV, calculated earlier in [12], see Table 4 of [14], and with wave functions fitted to the realistic wave functions in [16], (set I), while in set II parameters of the  $\Upsilon(nS)$  wave function were changed by 10-15%.

Results of calculations are given in Table 3, where we have put  $M_{br} \approx f_n$  and  $\left( \frac{M_\omega}{2\omega} \right)^2 = \frac{1}{2}$  to explain the decay  $\Gamma_{BB}(4S) = \Gamma_{exp}$  (see [16] for details).

**Table 3.** Values of  $\Gamma_\eta(n, n')$  (in keV) calculated using Eq. (12) with  $\left( \frac{M_{br}}{f_n} \frac{M_\omega}{2\omega} \right)^2 = \frac{1}{2}$  vs experimental data  $\Gamma_\eta^{exp}(n, n')$  (in keV).

| $(n, n')$              | (2,1)                                | (3,1)                        | (4,1)                 | (5,1) |
|------------------------|--------------------------------------|------------------------------|-----------------------|-------|
| $\Gamma_\eta$ , set I  | $2.50 \cdot 10^{-2}$                 | 1.45                         | 0.9                   | 3.5   |
| $\Gamma_\eta$ , set II | $1.4 \cdot 10^{-2}$                  | $3.6 \cdot 10^{-3}$          | 1.9                   | 1.3   |
| $\Gamma_\eta^{exp}$    | $(0.8 \pm 0.3) \cdot 10^{-2}$<br>[4] | $< 5.8 \cdot 10^{-3}$<br>[5] | $4.0 \pm 0.6$<br>[13] | -     |

Looking at the Table 3, one can see, that there is an order of magnitude agreement with experiment for the set II. Indeed, the factor  $\left(\frac{M_{br}}{f_\pi}\right)^2 \left(\frac{M_\omega}{2\omega}\right)^2$  can be estimated from  $\Upsilon(n, n')\pi\pi$  transitions studied in [14]-[18] to be roughly in the range  $[\frac{1}{2}, 2]$ . On theoretical side our formulas (7)-(11) automatically produce the width  $\Gamma_\eta(n, n')$  of the order of  $O(1 \text{ keV})$ , for all  $(n, 1)$  transitions except for  $(2, 1)$ , where a small phase space factor  $k^3$  gives two orders of magnitude suppression of  $\Gamma_\eta(2, 1)$ , and for  $\Gamma_\eta(3, 1)$ , which is highly sensitive to the form of wave function (cf. set I and set II). For the  $\Gamma_\eta(5, 1)$  one obtains a 7 keV value, which is however small as compared with the  $\Gamma_{\pi\pi}(5, 1)$ , the latter being  $O(1 \text{ MeV})$ . For  $\Gamma_\eta(4, 1)$  and  $\Gamma_{\pi\pi}(4, 1)$  from [16] the calculated ratio is  $R_{\eta/\pi\pi} \equiv \frac{\Gamma_\eta(4,1)}{\Gamma_{\pi\pi}(4,1)} \cong 3 \left(\frac{M_\omega}{2\omega}\right)^2 \left(\frac{f_\pi}{M_{br}}\right)^2 \approx 1.5$ , which roughly agrees with experimental value  $R_{\eta/\pi\pi}^{\text{exp}} = 2.41 \pm 0.40 \pm 0.12$  [13].

To compare our results with MEM predictions and experiment, we list in Table 4 the ratios  $\frac{\Gamma(\Upsilon(n,1)\eta)}{\Gamma(\psi(2,1)\eta)} \equiv X$  for  $n = 2, 3, 4, 5$  (the MEM numbers for  $n = 4, 5$  are obtained from  $n = 3$  using scaling  $X = O(p^3)$  [6].

**Table 4.** Values of the ratio  $\frac{\Gamma(\Upsilon(n,1)\eta)}{\Gamma(\psi(2,1)\eta)} \equiv X$  for  $n = 2, 3, 4, 5$  from MEM [6], experiment and the present paper (for the latter we take  $\Gamma_{\text{exp}}(\psi(2,1)\eta)$  in the denominator and two sets from Table 3):

| $n$           | 2                                    | 3                                | 4                       | 5                    |
|---------------|--------------------------------------|----------------------------------|-------------------------|----------------------|
| MEM           | $0.25 \cdot 10^{-2}$                 | $0.13 \cdot 10^{-2}$             | $0.33 \cdot 10^{-2}$    | $0.73 \cdot 10^{-2}$ |
| exper.        | $(0.8 \pm 0.2) \cdot 10^{-3}$<br>[4] | $< 0.58 \cdot 10^{-3}$<br>[5]    | $0.40 \pm 0.06$<br>[13] | -                    |
| present paper | $(1.4 \div 2.5) \cdot 10^{-3}$       | $(0.36 \cdot 10^{-3} \div 0.15)$ | $0.9 \div 1.9$          | $0.13 \div 0.35$     |

To check stability of our results, we have used for the wave function of  $B_s$  the realistic wave function different from that of  $B$ . As a result one obtains for  $\frac{\Gamma_\eta(n,1)}{\left(\frac{M_{br}}{f_\pi}\right)^2 \left(\frac{M_\omega}{2\omega}\right)^2}$  the values  $(2.74 \cdot 10^{-2}; 1.13; 0.44; 7.3) \text{ keV}$  for  $n = 2, 3, 4, 5$  respectively. To understand why  $\Gamma_\eta(.1)$  is so sensitive to the form of the wave function (cf. results for the sets I and II), we have varied the shape of wave functions of  $\Upsilon(nS)$  with  $n = 2, 3, 4, 5$  (explicitly the parameter  $\beta_1$ ) in the range  $\xi = \frac{\beta_1(n)}{\beta_1^{\text{opt}}(n)} = 0.8 \div 1.2$  of the optimal value, reproducing the realistic wave function. We have obtained the dependence of  $\Gamma_\eta(n, 1)$  on  $\xi$  for  $n = 2, 3, 4, 5$ . As a result  $\Gamma_\eta(2, 1)$  is rather stable, whereas  $\Gamma_\eta(3, 1)$  has minimum (almost zero) for  $\xi = 1.17$ . This fact explains results of sets I and II; while set I produces normal (and nearly maximal) values

for  $\Gamma_\eta(n, 1)$  of the order of 1 keV for  $n > 2$ , in case of set II the parameter  $\beta_1$  of  $\Upsilon(3S)$  roughly corresponds to the minimum of  $\Gamma_\eta$ . In this way doing comparison with experiment one obtains possibly an instrument for a precision study of wave functions of excited bottomonium states.

Summarizing, we have calculated the single  $\eta$  production width  $\Gamma_\eta(n, n')$  for  $\Upsilon(n, 1)\eta$  transitions with  $n = 2, 3, 4, 5$ . We have found that  $\Gamma_\eta(4, 1)$  are of the order of and larger than  $\Gamma_{\pi\pi}(4, 1)$ . This fact is in agreement with the latest measurements in [13] of  $\Gamma_\eta^{\text{exp}}(4, 1)$ . We have shown that  $\Gamma_\eta(n, 1)$ ,  $n = 4, 5$  is large ( $\sim O(1 \text{ keV})$ ) for typical (realistic) parameters of  $\Upsilon(nS)$  wave functions, but can occasionally drop near zero for slightly varied form of wave function, as it happens for  $n = 3$ . This high sensitivity is connected to the oscillating character of excited bottomonium wave functions<sup>2</sup>. Our calculations do not contain fitting parameters; the only two parameters  $M_{br}, M_\omega$  are fixed by previous comparison with dipion data. One should stress that  $\eta$  production in bottomonium is not suppressed in our approach as compared to  $\eta$  production in charmonium transitions. This is in contrast with the results of method of [6, 7]. We have given arguments why the dipion transitions in high excited states of heavy quarkonia as well as single  $\eta$  and  $\pi$  emission cannot be reliably calculated within the MEM method of [6, 7], widely used now. As it is seen in Tables 3 and 4 the sequence of experimental data [4],[5], [13] contradict predictions of MEM. Recently a new calculation was done of  $\Gamma_\eta(n, n')$  in [20] where also  $B\bar{B}^*$  etc. intermediate states were taken into account as well as in our approach. The authors however did not use wave functions of hadrons involved, but rather exploited fitted coupling constants and formfactors, and specific form of matrix elements, which makes it difficult to compare with our method directly.

The authors are grateful to S.I.Eidelman for constant support and helpful comments.

The financial support of grants RFFI 06-02-17012, 06-02-17120 and NSh-4961.2008.2 is gratefully acknowledged.

## References

- [1] W.-M. Yao et al, Particle Data Group, J. Phys., G **32**(2006) 1.
- [2] V.Fonseca et al., (CUSB Collaboration), Nucl. Phys., B **242**(1984) 31.
- [3] I.C.Brock et al., Phys. Rev., D **43** (1991) 1448.
- [4] D.Kreinick et al., (CLEO Collaboration), arXiv: 0710.5929.

---

<sup>2</sup>A similar phenomenon of cancellation in matrix elements was mentioned in [6]

- [5] Q.He et al, (CLEO collaboration), arXiv:0806.3027.
- [6] T.-M.Yan, Phys. Rev., D **22** (1980) 1632;  
Y.-P.Kuang, Front. Phys. China, **1**(2006) 19 , arXiv: hep-ph/0601044.
- [7] M.B.Voloshin, arXiv: 0711.4556; M.B.Voloshin, V.I.Zakharov, Phys. Rev. Lett. **45** (1980) 688.
- [8] M.Campostrini, A.Di Giacomo and G.Mussardo, Z.Phys., C **25** (1984) 173;  
A.Di Giacomo, and S.Olejnik, Z. Phys., C **34** (1986) 577; G.Bali, N. Brambilla, A.Vairo, Phys., Lett. B **421** (1998) 265.
- [9] for a review see A.Di Giacomo, H.G.Dosch, V.I.Shevchenko, Yu.A.Simonov, Phys. Rept. **372** (2002) 319.
- [10] H. Leutwyler, Phys. Lett., **98B** (1981) 447;  
M. B. Voloshin, Nucl. Phys., B **154** (1979) 365; Sov. J. Nucl. Phys., **36** (1982) 143.
- [11] Yu.A.Simonov, S.Titard, F.J.Yandurain, Phys. Lett. B **354** (1995) 435.
- [12] A.M.Badalian, A.I.Veselov, B.L.G.Bakker, J.Phys. G **31**(2005) 417  
Phys. Atom Nucl. **70** (2007) 1764 .
- [13] N.Arnaud (for BaBar Collaboration), arXiv: 0805.2591.
- [14] Yu.A.Simonov, Phys. Atom. Nucl. **71** (2008) 1048, arXiv: 0711.3626.
- [15] Yu.A.Simonov, JETP Lett., **87** (2008) 147, arXiv: 0712.2197.
- [16] Yu.A.Simonov, A.I.Veselov, arXiv: 0804.4635; Phys. Rev. D (in press).
- [17] Yu.A.Simonov, A.I.Veselov, Phys. Lett. B **671** (2009) 55, arXiv:0805.4499.
- [18] Yu.A.Simonov, A.I.Veselov, JETP Lett. **88** (2008) 79, arXiv: 0805.4518.
- [19] Yu.A.Simonov, Phys.Rev. D **65** (2002) 0940018.
- [20] Ce Meng, K.-T.Chao, Phys. Rev. D **78** (2008) 074001.

## Appendix 1

### Matrix element of single $\eta$ emission

According to the general theory in [14, 16], the matrix elements  $\mathcal{M}_\eta^{(1)}, |Mc_\eta^{(2)}$  for  $\Upsilon(n, n')\eta$  corresponding to diagrams of Fig.1, (a) and (b) respectively, can be written as

$$\mathcal{M}_\eta^{(1)}(n) = \frac{M_{br}M_\omega}{f_\eta N_c \sqrt{2\omega_\eta}} \int \frac{d^3p}{(2\pi)^3} \sum_{n_2, n_3} \frac{J_{nn_2n_3}(\mathbf{p}, \mathbf{k}) J_{n'n_2n_3}^+(\mathbf{p})}{E - E_{n_2n_3}(\mathbf{p})}. \quad (\text{A.1})$$

Here  $n_2, n_3$  are channels of intermediate state, with e.g.  $n_2 = B, B^*, B_s, B_s^*, \dots$ , we omit indices  $n_2, n_3$  and write

$$Jn(\mathbf{p}, \mathbf{k}) = \int \bar{y}_{n23}^{(\eta)} \frac{d^3\mathbf{q}}{(2\pi)^3} \tilde{\Psi}_n(c\mathbf{p} - \frac{\mathbf{k}}{2} + \mathbf{q}) \tilde{\psi}_{n_2}(\mathbf{q}) \tilde{\psi}_{n_3}(\mathbf{q} - \mathbf{k}) \quad (\text{A.2})$$

where  $\tilde{\Psi}_n, \tilde{\psi}_{n_i}$  are momentum space wave functions of  $\Upsilon(nS)$  and  $B(B^*)$  mesons respectively.

The vertex factor  $\bar{y}_{123}^{(\eta)}$  is calculated in the same way as in [14], namely from the Dirac trace of the projection operators for the decay process in our case this is  $\Upsilon(nS) \rightarrow BB^*\eta$ . Identifying the creation operators as  $\bar{\psi}_b\gamma_i\psi_b, \bar{\psi}_b\gamma_5\psi_n, \bar{\psi}_b\gamma_k\psi_n$ ,  $n = u, d, s$  and extracting vertex of  $\eta$  creation from the Lagrangian  $\Delta\mathcal{L} = -\int \bar{\psi}_n \hat{U} M_{br} \psi_n d^4x$

which gives  $i \frac{M_{br} \bar{\psi}_n \gamma_5 \hat{\lambda} \psi_n}{f_n \sqrt{2\omega_\eta}}$ , with  $\hat{\lambda} = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & & \\ & 1 & \\ & & -2 \end{pmatrix}$ , one has for the decay process (cf. Appendix 1 of [14])

$$G(\Upsilon \rightarrow BB^*\eta) = \text{tr}[\gamma_i S_b(u, w) \gamma_5 S_{\bar{n}}(w, x) \gamma_5 S_n(x, w) \gamma_k S_{\bar{b}}(w', v)] \quad (\text{A.3})$$

As shown in [14], appendix 1 and 2, the quark Green's functions can be split into two factors  $S(x, y) = \Lambda^\pm G(x, y)$ , with the projection operators  $\Lambda_k^\pm = \frac{m_k \pm \omega_k \gamma_4 \mp i p_i^{(k)} \gamma_i}{2\omega_k}$ ,  $k = b, n$  and the scalar part  $G(x, y)$ , where spins are present only in spin-dependent interaction and treated as corrections. Here  $\omega_k$  is the average energy of quark in given meson. Hence one is brought to the spin factor  $Z$ .

$$Z = \text{tr}(\gamma_i \Lambda_b^+ \gamma_5 \Lambda_n^- \gamma_5 \Lambda_n^+ \gamma_k \Lambda_b^-) \quad (\text{A.4})$$

which is equal to

$$Z = \frac{m_b^2 + \Omega^2}{4\omega^2 \omega^2} ((\omega^2 - \mathbf{p}^q \mathbf{p}^{\bar{q}}) \delta_{ik} - p_i^q p_k^{\bar{q}} + p_k^q p_i^{\bar{q}}). \quad (\text{A.5})$$

Here  $\Omega, \omega$  are average energies of  $b$  and  $n$  quark in  $B$  or  $B^*$ . One can identify the momenta of  $B$  and  $B^*$  as  $\mathbf{P}_1 = \mathbf{p}$  and  $\mathbf{P}_2 = -\mathbf{p} - \mathbf{k}$ , then  $\mathbf{q}$  in (A.2) can be expressed as

$$\mathbf{p}_{\bar{q}} = -\mathbf{q} + \frac{\omega}{\omega + \Omega} \mathbf{p}, \quad \mathbf{p}_q = \mathbf{q} - \frac{\omega}{\omega + \Omega} \mathbf{p} - \mathbf{k} \frac{\Omega + 2\omega}{\Omega + \omega}, \quad (\text{A.6})$$

and  $Z$  is (we put  $m_b \cong \Omega$ )

$$Z = \frac{1}{2\omega^2}(-\mathbf{k}\mathbf{q}\delta_{ik} + k_i q_k + k_k q_i). \quad (\text{A.7})$$

It is important, that we are looking for the  $P$ -wave of emitted  $\eta$ , and hence for  $P$  wave of relative  $BB^*$  motion, hence the integral (A.2) should yield the term  $\mathbf{p}\mathbf{k}$ . This indeed happens, when one approximates  $\tilde{\Psi}_n, \tilde{\psi}_n$  as series of oscillator wave functions and (A.2) has the form

$$I_n(\mathbf{p}, \mathbf{k}) = \bar{y}_{n23}^\eta e^{-\frac{\mathbf{p}^2}{\Delta_n} - \frac{\mathbf{k}^2}{4\beta_2^2}} {}^{(0)}I_n(\mathbf{p}). \quad (\text{A.8})$$

In the process of  $d\mathbf{q}$  integration in (A.2) one changes the integration variable  $q_i \rightarrow q'_i - u_n + O(k_i)$  with  $\beta_2, \Delta_n$  are oscillator parameters, found by  $chi^2$  procedure.

Thus result of  $d^3q$  integration yields

$$\bar{y}_{123}^\eta = \frac{u_n}{2\omega^2}(-\mathbf{k}\mathbf{p}\delta_{ik} - k_i p_k + K_k p_i). \quad (\text{A.9})$$

In an analogous way one obtains for  $J_{n'}(\mathbf{p})$  in (A.1) the form

$$J_{n'}(\mathbf{p}) = \bar{y}_{n'23}^{(\eta)} e^{-\frac{\mathbf{p}^2}{\Delta_{n'}}} {}^{(1)}I_{n'}(\mathbf{p}) \quad (\text{A.10})$$

and  $\bar{y}_{n'23}^{(\eta)}$  is obtained from the Dirac trace for the process  $B\bar{B}^* \rightarrow \Upsilon(n'S')$ ,

$$Z_2(B\bar{B}^*) = \frac{1}{2\omega} e_{i'kl}(-2q_l + \frac{2\omega}{\omega + \Omega} p_l) \quad (\text{A.11})$$

and the result of integration over  $d^3q$  yields in (A.10)

$$\bar{y}_{n'23}^{(\eta)} = -e_{i'kl} \frac{p_l}{\omega}. \quad (\text{A.12})$$

Here  $i'$  is the polarization of  $\Upsilon(n'S)$  (represented by  $\bar{\psi}_b \gamma_{i'} \psi_b$ ) and  $k$  as in (A.9) is the polarization of  $B^*$ . Averaging over angles of  $\mathbf{p}$  one obtains

$$\langle \bar{y}_{n23}^{(\eta)} \bar{y}_{n'23}^{(\eta)} \rangle_p = \frac{u_n}{3} \frac{\mathbf{p}^2}{\omega^3} (e_{i'il} k_l) \quad (\text{A.13})$$

and finally one writes as in (9)

$$\mathcal{M}_n^{(1)}(n, n') = \frac{M_\omega M_{br} u_n 2e_{i'il} k_l}{f_n \sqrt{2\omega_\eta} \sqrt{3} \cdot 3} \left( \frac{\mathcal{L}^{(1)}}{\omega^3} - \frac{\mathcal{L}_s^{(1)}}{\omega_s^3} \right) e^{-\frac{k^2}{4\beta_2^2}}. \quad (\text{A.14})$$

For  $\mathcal{M}_n^{(2)}(n, n')$  one can use time inversion and interchange indices  $i, i'$  and change sign of  $\mathbf{k}$ , obtaining in this way Eqs. (9) and (10) of the main text. For the intermediate state of  $B^* \bar{B}^*$  the summation over polarizations of  $B^*$  yields a net zero result, therefore we are left with only  $(B\bar{B}^* + B^* \bar{B})$  intermediate state.